

Association between Four Measures of Retention in Care and Virologic Failure

Kalmin MM, Castel AD, Greenberg AE, Young H, Hoffman HJ, on Behalf of the DC Cohort Executive Committee
The George Washington University, Milken Institute School of Public Health

BACKGROUND

- Sustaining viral suppression (VS) is the ultimate goal of HIV care and treatment.
- Retention in care is a critical step along the HIV care continuum leading to VS.
- Retention has been measured in various ways without identification of a gold standard for predicting VS.
- Therefore, the relationship between various retention measures and sustained VS merits further exploration.

OBJECTIVES

- To compare the association between four retention measures and virologic failure (VF) among an urban cohort of HIV-infected persons in care

METHODS

DC COHORT

- A longitudinal observational study of HIV-infected persons in care in Washington, DC at 15 clinical sites
- Data abstracted from participants' medical records and linked to DC Dept. of Health data
- Included participants enrolled for at least 12 months from 1/2011 to 9/2014, age ≥ 18 years from 9 sites, with an undetectable viral load at study entry

ANALYSIS

- Outcome: VF defined as a viral load (VL) >200 copies/ml during follow-up
- Exposure: retention as measured in four ways
 - No gaps in care > 6 months
 - 4-month visit constancy
 - Institute of Medicine (IOM) measure: >2 visits at least 90 days apart in 12-months
 - Health and Human Services (HHS) measure: >1 visit in each 6-month interval in 24 months with ≥ 60 days between visits
- Performed multivariable Cox proportional hazards regression to assess the relationship between each retention measure and VF, modeled as a recurrent event

RESULTS

Table 1. Socio-Demographics, Clinical Characteristics, and Retention Estimates of HIV-Infected Adult Patients with an Undetectable Viral Load at Enrollment in the DC Cohort (N=1,958)

Characteristic	Median (IQR) or Frequency (%)
Median study follow up time (months)	35.6 (27.6-40.0)
Median baseline age (years)	49.3 (41.0-56.2)
Sex at birth	
Male	1,530 (78.1%)
Female	428 (21.9%)
Race/ethnicity ¹	
Non-Hispanic black	1,402 (71.6%)
Non-Hispanic white	420 (21.5%)
Other/unknown	136 (6.9%)
Insurance type ¹	
Private	681 (34.8%)
Medicaid	449 (22.9%)
Medicare	286 (14.6%)
Other/unknown	542 (27.7%)
HIV transmission risk ¹	
MSM ²	1,055 (53.9%)
High risk heterosexual	527 (26.9%)
IDU	267 (13.6%)
Other/unknown	109 (5.6%)
DC Cohort site type	
Academic/hospital based organization	1195 (61.0%)
Community based organization	763 (39.0%)
Median duration of HIV infection at baseline (years) ¹	11.9 (6.5-23.1)
Median baseline CD4 cell count (cells/ μ l) ¹	568 (407-763)
Retention Estimates ³	
Median % time without a 6-month gap in care	100 (79.7-100)
Median 4-month visit constancy	66.7 (50-87.5)
Median IOM continuous care measure	66.7 (50-100)
Median HHS retention in medical care ⁴	50 (0-100)

¹ Variables for which the DC DOH may have contributed data

² Includes patients who reported both MSM and IDU transmission risk

³ Retention estimates reflect percentage of follow up time in which patients met the specific measure

⁴ Limited to only those patients with at least 24 months of follow up (n=1,617)

Table 2. Associations between Measures of Retention in HIV Care and Time to Virologic Failure among Patients with Undetectable Viral Load at Study Enrollment





	Retention Measure ¹	Follow-up time	Hazard ratio (95% CI)
Analysis #1: Patients with at least 12 months of follow up (N=1,958) ²	% of time without a gap > 6 months	0-2 yrs	1.24 (1.06-1.45)
		2-4 yrs	0.896 (0.676-1.19)
	% of time meeting 4-month visit constancy	0-2 yrs	1.03 (0.912-1.17)
		2-4 yrs	0.84 (0.658-1.07)
Analysis #2: Patients with at least 24 months of follow up (N=1,617) ²	% of time meeting IOM continuous care	0-2 yrs	0.97 (0.880-1.07)
		2-4 yrs	0.74 (0.615-0.885)
	% of time meeting HHS retention in medical care	0-2 yrs	0.97 (0.902-1.05)
		2-4 yrs	0.87 (0.761-0.982) ³

¹ Retention measures modeled as per 25% increase in retention

² Adjusted for baseline age, sex, race/ethnicity, insurance type, HIV transmission risk, site type, duration of follow up, baseline CD4 count, baseline ART regimen type, change in ART regimen during follow up, and baseline AIDS status

³ Not statistically significant under the Bonferroni corrected p-value for multiple comparisons

Figures 1- 4. Kaplan Meier Curves of Virologic Failure Comparing Quartiles for Each Retention in Care Measure



CONCLUSIONS

SUMMARY

- Both crude (Figures 1- 4) and adjusted associations (Table 2) between retention and VF varied by retention measure.
- During the first 2 years of follow-up, an increasing percent of time without a gap in care >6 months was associated with an increased rate of VF (aHR: 1.24, 95% CI: 1.06-1.45).
- During the last 2 years of follow-up, an increasing percent of time in which the IOM measure was met was associated with a decreased rate of VF (aHR: 0.74, 95% CI: 0.62-0.89).
- Other retention measures were not associated with time to VF.

LIMITATIONS

- Data reflect only those patients who consented to be in the DC Cohort.
- VL data were limited to consent date forward, so there is a lack of information on patients' histories of VS prior to study enrollment.
- Retention may have been underestimated as VL tests could not be used as proxies for routine HIV care visits since values of VLs were also used to estimate VF.

DISCUSSION

- Our results suggest that monitoring retention and its association with VF varies depending on how retention is defined.
- After 2 years, the rate of VF declined when the IOM measure was met.
- None of the retention measures was associated with a decreased rate of VF during the first 2 years of follow-up.
- Early in follow-up, there was a direct relationship between the no gaps in care > 6 months measure and VF, a finding that warrants additional analysis.
- Further follow-up will enable the assessment of whether these patterns are maintained over time.

For additional information please contact: Mariah M. Kalmin at mariahm@gwu.edu